EARLY DETECTION IS KEY

Everyone should get tested for HBV infection because hepatitis B is one of the greatest health threats for Hmong.

» Get Tested

Ask your doctor for these blood tests:

Hepatitis B surface antigen (HBsAg):

Tells if you have chronic hepatitis B (also known as being a hepatitis B carrier).

Hepatitis B surface antibody (anti-HBs):

Tells if you are protected against HBV.

Only the HBsAg blood test can tell if you have chronic hepatitis B.



» Get Vaccinated

If both your HBsAg and anti-HBs blood tests are negative, then you are not protected and should get vaccinated. Make sure your baby also gets vaccinated at birth.

The hepatitis B vaccine is a 3 shot series given over 6 months that can protect you for life.

The vaccine is so effective at preventing HBV and liver cancer that it is called "the first anti-cancer vaccine" by the World Health Organization.

Unite against HBV.



Jade represents good health and longevity.
The **Jade Ribbon** spreads awareness and supports
people worldwide living with chronic
hepatitis B and liver cancer.





Asian Liver Center at Stanford University

780 Welch Road, CJ 130
Palo Alto, CA 94304
Tel: 888.311.3331
Fax: 650.736.8001
liver.stanford.edu
#IOINJADE

ENGLISH-HMONG EDITION
© 2016 ASIAN LIVER CENTER





BREAKING THE SILENCE ABOUT HBV

- » Hepatitis B virus (HBV) is the most common serious infection of the liver, and can lead to a shortened lifespan from liver cancer or liver failure.
- » Up to 1 in 6 Hmong are living with chronic (life-long) hepatitis B.



- » Most don't even know that they're infected.
- » 1 in 4 people with chronic hepatitis B will die from liver cancer or liver failure, but this can be avoided with appropriate monitoring and treatment.

EVEN IF YOU FEEL HEALTHY, YOU MAY ALREADY BE INFECTED

- » Hepatitis B is dangerous because there are often no symptoms.
- » By the time symptoms such as stomach pain or jaundice (dark urine and yellowing of the skin or eyes) appear, it is often too late for treatment to be effective.

TRANSMITTED JUST LIKE HIV



1. A mother-to-child infection For Hmong, HBV is commonly

transmitted from an infected mother to her child at birth.

2. A blood-borne infection, for example:

- » Wound-to-wound contact
- » Sharing razors or toothbrushes
- » Reusing needles for tattoos, piercings, or injecting drugs
- » Reusing medical needles or syringes



3. A sexually transmitted infection

HBV can be transmitted through unprotected sex with an infected person.

HBV IS NOT TRANSMITTED THROUGH FOOD AND WATER

It is important to understand that HBV is **NOT** transmitted through:

- » Sharing food and water
- » Sharing utensils or drinking glasses
- » Casual contact
- » Coughing or sneezing
- » Hugging or kissing
- » Breastfeeding

There is no reason to distance yourself from those infected with HBV.

If you have chronic hepatitis B, you should not be excluded from work, school, or other daily activities.

WHAT YOU SHOULD DO IF YOU HAVE HBV

» Get screened

EVEN THOUGH YOU MAY FEEL HEALTHY, REGULAR SCREENING IS IMPORTANT.

Ask your doctor for the following tests:

Every	Test	Screens for
6 months	ALT blood test	liver damage
	AFP blood test	liver cancer
1 year*	ultrasound	liver cancer

* If you have liver cirrhosis or your family has a history of liver cancer, increase the frequency of your ultrasound exam to every 6 months.

If detected at an early stage, liver cancer can be successfully treated.

» Not everyone who tests positive for HBV needs treatment

But if your blood ALT is high, this means you have liver damage. So, you may start to take anti-viral medicine to prevent HBV from causing more damage. Even though there is no cure for hepatitis B, appropriate treatment can reduce your risk of liver cancer and liver failure. Ask your doctor about FDA-approved treatment for HBV.

» Get the hepatitis A vaccine

- » Avoid drinking alcohol
- » Protect your loved ones

Make sure your family and partner are tested for HBV and vaccinated if they are not protected.

If you are pregnant, make sure your newborn receives the first shot of the hepatitis B vaccine **and** the hepatitis B immunoglobulin (HBIG) shot at birth. Then complete the hepatitis B vaccine series on time. This will protect your newborn from infection.